Facts about UNDESCENDED TESTICLE

UNDESCENDED TESTICLE IS ALSO CALLED CRYPTORCHIDISM

PAINLESS
CONDITION

CRYPTORCHIDISM IS
CONGENITAL
(MEANING IT IS
PRESENT AT BIRTH)



IT IS FOUND IN ABOUT



3-4% OF NEWBORN BOYS

AND UP TO 21% OF PREEMIE BOYS

10% OF CASES INVOLVE BOTH TESTES



Facts about Undescended Testicle



American Urological Association

IN MOST CHILDREN

WITH THIS HEALTH ISSUE, IT'S NOT KNOWN WHY THE **TESTICLES FAIL TO DROP**



NOT SEEING OR FEELING A

TESTICLE WHERE YOU WOULD EXPECT IT TO BE IN THE SCROTUM IS THE

UNDESCENDED TESTICLE

MYTH: AN UNDESCENDED TESTICLE IS ALWAYS **OBVIOUS** IN BABIES

ABOUT 20 PERCENT OF CAS-ES ARE FOUND ONCE A BOY IS NO LONGER A BABY. ALL BOYS SHOULD HAVE THE LOCATION OF THEIR TESTICLES CHECKED DURING EACH ANNUAL PHYSICAL EXAM.

— even those whose testicles have properly descended — should learn how to do a testicular self-exam when they are teenagers to detect any lumps or bumps that might be early signs of medical problems

ABOUT

OF UNDESCENDED **TESTICLES DROP**

ON THEIR **OWN** DURING THE

FIRST THREE







OF MALES WITH AN UNDESCENDED **TESTICLE ALSO**

HAD A MALE **RELATIVE WITH** THE CONDITION



IN MOST CASES, BOYS WITH ONE UNDESCENDED

TESTICLE ARE STILL ABLE TO HAVE CHILDREN OF THEIR OWN **LATER IN LIFE**